




Please replace formula IIIaf on page 28 with the following rewritten formula:

C<sup>17</sup>


Please replace formula IVo on page 32 with the following rewritten formula:

C<sup>18</sup>

Please replace formula Vi on page 34 with the following rewritten formula:

C<sup>19</sup>


Please replace formula Vt on page 36 with the following rewritten formula:

C<sup>20</sup>

### IN THE CLAIMS:

Please amend claims 11, 13, 14, 33, 34, 36, and 38, and enter new claims 41 and 42, as follows:

- Sub  
D1/ 11. (Amended) A method for providing sedation, mitigating anxiety or providing anaesthesia in a subject in need thereof, comprising administering to a subject an effective amount of a therapeutic compound, wherein the therapeutic compound is of the formula (Ia):



C21


in which  $F^2$  is an organic radical which may be joined in a cyclic ring system with  $G^2$ , and which may contain inorganic counterions, but is not a nitrate group; E is a methylene group and  $G^1$  is a methylene group or does not exist;  $F^1$  is H; and  $G^2$  is  $R^N \cdot Z^N \cdot$ ;

wherein  $R^N$  is an organic radical possessing a heteroaryl group containing P or S atoms where said P or S are positioned  $\beta$ ,  $\gamma$ , or  $\delta$  to a nitrate group as identified in formula I; and  $Z^N$  is  $W^N_{mm} \cdot X^N_{nn} \cdot Y^N_{oo}$ ;

wherein mm, nn, oo are 0 or 1 and  $W^N, X^N, Y^N$  are  $NH, NR^{NN}, CO, O$  or  $CH_2$ ;

wherein  $R^{NN}$  is a short chain alkyl group ( $C_1 - C_{12}$ ).

- Sub  
D2/ 13. (Twice Amended) A method for providing sedation, mitigating anxiety or providing anaesthesia in a subject in need thereof, comprising administering to a subject an effective amount of a therapeutic compound, wherein the therapeutic compound is of the formula (Ic):



C22

in which E is  $(R^1 R^2 C)_m$  and  $G^2 - G^1 - CF^1 F^2 -$  is  $R^{19} - (R^3 R^4 C)_p - (R^{17} R^{18} C)_n -$ ;

wherein: m, n, p are integers from 0 to 10;

$R^{3,17}$  are each independently hydrogen, a nitrate group, or A; and

$R^{1,4}$  are each independently hydrogen, or A;

Sub  
P<sub>2</sub>  
cont

C22  
cont.

where A is selected from a substituted or unsubstituted aliphatic group (comprising a branched or straight-chain aliphatic moiety having from 1 to 24 carbon atoms in the chain, which optionally may contain O, S, NR<sup>6</sup> and unsaturations in the chain, optionally bearing from 1 to 4 hydroxy, nitrate, amino, aryl, or heterocyclic groups; an unsubstituted or substituted cyclic aliphatic moiety having from 3 to 7 carbon atoms in the aliphatic ring, which optionally may contain O, S, NR<sup>6</sup> and unsaturations in the ring, optionally bearing from 1 to 4 hydroxy, nitrate, amino, aryl, or heterocyclic groups; an unsubstituted or substituted aliphatic moiety constituting a linkage of from 0 to 5 carbons, between R<sup>1</sup> and R<sup>3</sup> and/or between R<sup>17</sup> and R<sup>4</sup>, which optionally may contain O, S, NR<sup>6</sup> and unsaturations in the linkage, and optionally bearing from 1 to 4 hydroxy, nitrate, amino, aryl, or heterocyclic groups); a substituted or unsubstituted aliphatic group (comprising a branched, cyclic or straight-chain aliphatic moiety having from 1 to 24 carbon atoms in the chain) containing carbonyl linkages (C=O, C=S, C=NOH), which optionally may contain O, S, NR<sup>6</sup> and unsaturations in the chain, optionally bearing from 1 to 4 hydroxy, nitrate, amino, aryl, or heterocyclic groups; a substituted or unsubstituted aryl group; a heterocyclic group; an amino group selected from alkylamino, dialkylamino, cyclic amino, diamino and triamino moieties, arylamino, diarylamino, and alkylarylamino; hydroxy; alkoxy; a substituted or unsubstituted aryloxy;

wherein X is F, Br, Cl, NO<sub>2</sub>, CH<sub>3</sub>, CF<sub>2</sub>, O, NH, NMe, CN, NHOH, N<sub>2</sub>H<sub>3</sub>, N<sub>2</sub>H<sub>2</sub>R<sup>13</sup>, N<sub>2</sub>HR<sup>13</sup>R<sup>14</sup>, N<sub>3</sub>, S, SCN, SCN<sub>2</sub>H<sub>2</sub>(R<sup>15</sup>)<sub>2</sub>, SCN<sub>2</sub>H<sub>3</sub>(R<sup>15</sup>), SC(O)N(R<sup>15</sup>)<sub>2</sub>, SC(O)NHR<sup>15</sup>, SO<sub>3</sub>M, SH, SR<sup>7</sup>, SO<sub>2</sub>M, S(O)R<sup>8</sup>, S(O)<sub>2</sub>R<sup>9</sup>, S(O)OR<sup>8</sup>, S(O)<sub>2</sub>OR<sup>9</sup>, PO<sub>2</sub>HM, PO<sub>3</sub>HM, PO<sub>3</sub>M<sub>2</sub>, P(O)(OR<sup>15</sup>)(OR<sup>16</sup>), P(O)(OR<sup>16</sup>)(OM), P(O)(R<sup>15</sup>)(OR<sup>8</sup>), P(O)(OM)R<sup>15</sup>, CO<sub>2</sub>M, CO<sub>2</sub>H, CO<sub>2</sub>R<sup>11</sup>, C(O), C(O)R<sup>12</sup>, C(O)(OR<sup>13</sup>), PO<sub>2</sub>H, PO<sub>2</sub>M, P(O)(OR<sup>14</sup>), P(O)(R<sup>13</sup>), SO, SO<sub>2</sub>, C(O)(SR<sup>13</sup>), SR<sup>5</sup>, SSR<sup>7</sup> or SSR<sup>5</sup>;

Y is F, Br, Cl, CH<sub>3</sub>, CF<sub>2</sub>H, CF<sub>3</sub>, OH, NH<sub>2</sub>, NHR<sup>6</sup>, NR<sup>6</sup>R<sup>7</sup>, CN, NHOH, N<sub>2</sub>H<sub>3</sub>, N<sub>2</sub>H<sub>2</sub>R<sup>13</sup>, N<sub>2</sub>HR<sup>13</sup>R<sup>14</sup>, N<sub>3</sub>, S, SCN, SCN<sub>2</sub>H<sub>2</sub>(R<sup>15</sup>)<sub>2</sub>, SCN<sub>2</sub>H<sub>3</sub>(R<sup>15</sup>), SC(O)N(R<sup>15</sup>)<sub>2</sub>, SC(O)NHR<sup>15</sup>, SO<sub>3</sub>M, SH, SR<sup>7</sup>, SO<sub>2</sub>M, S(O)R<sup>8</sup>, S(O)<sub>2</sub>R<sup>9</sup>, S(O)OR<sup>8</sup>, S(O)<sub>2</sub>OR<sup>9</sup>, PO<sub>2</sub>HM, PO<sub>3</sub>M<sub>2</sub>, P(O)(OR<sup>15</sup>)(OR<sup>16</sup>), P(O)(OR<sup>16</sup>)(OM), P(O)(R<sup>15</sup>)(OR<sup>8</sup>), P(O)(OM)R<sup>15</sup>, CO<sub>2</sub>M, CO<sub>2</sub>H, CO<sub>2</sub>R<sup>11</sup>, C(O)R<sup>12</sup>, C(O)(OR<sup>13</sup>), C(O)(SR<sup>13</sup>), SR<sup>5</sup>, SSR<sup>7</sup> or SSR<sup>5</sup>, or does not exist;

R<sup>2</sup>, R<sup>5</sup>, R<sup>18</sup>, R<sup>19</sup> are optionally hydrogen, A or X-Y;

R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup> are the same or different alkyl or acyl groups containing 1-24 carbon atoms which may contain 1-4 ONO<sub>2</sub> substituents; or C<sub>1</sub> - C<sub>6</sub> connections to R<sup>1</sup> - R<sup>4</sup> in cyclic derivatives which may contain 1-4 ONO<sub>2</sub> substituents; or are each independently hydrogen, a nitrate group or A;

C 22  
cont.

M is H, Na<sup>+</sup>, K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, N<sup>+</sup>H<sub>k</sub>R<sup>11</sup> where k is 0-3; or other pharmaceutically acceptable counterion;

and with the proviso that when m = n = p = 1 and R<sup>19</sup>, R<sup>2</sup>, R<sup>18</sup>, R<sup>1</sup> = H and R<sup>17</sup>, R<sup>3</sup> are nitrate groups, R<sup>4</sup> is not H.

C 23  
cont.

14. (Amended) The method of claim 11, wherein F<sup>2</sup> is a nitrate group; and E, F<sup>1</sup>, G<sup>1</sup>, G<sup>2</sup> are the same or different organic radicals which may be joined in cyclic ring systems, and which may contain inorganic counterions;

with the proviso that when E and G<sup>1</sup> are methylene groups and F<sup>1</sup> is H, G<sup>2</sup> is not a nitrate group, nor R<sup>N</sup>Z<sup>N</sup>;

wherein R<sup>N</sup> is any aryl or heteroaryl group and Z<sup>N</sup> is (CO)<sub>mm</sub>-X<sup>N</sup><sub>nn</sub>-Y<sup>N</sup><sub>oo</sub>;


wherein mm, nn, oo are 0 or 1 and X<sup>N</sup>, Y<sup>N</sup> are NH, NR<sup>NN</sup>, O or CH<sub>2</sub>;

wherein R<sup>NN</sup> is a short chain alkyl group (C<sub>1</sub> - C<sub>12</sub>).


33. (Amended) A method of providing sedation or mitigating anxiety in a subject in need thereof, comprising administering to a subject an effective amount of a therapeutic compound selected from the group consisting of:

C 24


IIIa




IIIb




IIIc




IIIId




IIIe




IIIf




IIIg




IIIh




IIIi

24  
cont.




IIIj








IIIk









IIIl







C 24  
cont.III<sub>m</sub>III<sub>n</sub>



124  
cont.


III<sub>t</sub>III<sub>u</sub>III<sub>v</sub>III<sub>w</sub>III<sub>x</sub>III<sub>y</sub>24  
cont.






C24  
cont.


IIIaf




IIIag




IIIah




IIIai




IIIaj




IIIak








24  
cont.




34. (Amended) The method of claim 33, wherein the compound has the formula IIIc:




36. (Amended) A method of providing sedation or mitigating anxiety in a subject in need thereof, comprising administering to a subject an effective amount of a therapeutic compound selected from the group consisting of:




IVc




IVd




IVe




IVf









IVg








IVh


C 25  
Cont.




25  
cont.




38. (Amended) A method of mitigating anxiety in a subject in need thereof, comprising administering to a subject an effective amount of a therapeutic compound selected from the group consisting of:




Vb




Vc




Vd




Ve




Vf




Vg




Vh




Vi




Vj




Vk




Vl



Vm




Vn







Vo




Vp




Vq




Vr




Vs




Vt

26  
cont.


Vu




Vv




Vw




Vx




Vy




Vz

126  
cont.


Vaa




Vab




Vac




Vad




Vae



Vaf



Vag



, and